

# A GENERALIZED $\sigma$ -POROUS SET WITH A SMALL COMPLEMENT

JAROSLAV TIŠER

*Received 29 October 2003*

We show that in every Banach space, there is a  $\sigma$ -porous set, the complement of which is of  $\mathcal{H}^1$ -measure zero on every  $C^1$  curve.

## 1. Introduction

The notion of porosity is one of the notions of smallness which had appeared in a natural way in the theory of differentiation of functions. Porous and  $\sigma$ -porous sets play the role of the exceptional sets in many contexts. A survey as well as further references on this topic can be found in [4]. In finite-dimensional Banach spaces, the family of all  $\sigma$ -porous sets is a proper subfamily of the sets of both 1st category and Lebesgue-measure zero.

The question how big a porous or a  $\sigma$ -porous set can be in an infinite-dimensional Banach space is closely related to the existence of a point of Fréchet differentiability of a Lipschitz function. The simplest connection may be the easy observation that for a porous set  $M$ , the distance function  $f(x) = \text{dist}(x, M)$  is a Lipschitz function which is not Fréchet differentiable at any point of  $M$ . Another more important link is represented by a deep result in [1]. From this result it follows, in particular, that *every Lipschitz function on a separable Asplund space is Fréchet differentiable except for the points belonging to the union of a  $\sigma$ -porous set and a  $\Gamma$ -null set*. The concept of a  $\Gamma$ -null set is another and relatively new concept of smallness of a set. We will not give here the definition (it can be found in the just-mentioned paper [1]) since it will not be used here. We only remark that  $\Gamma$ -null sets form a  $\sigma$ -ideal of subsets and in a finite-dimensional space, the  $\Gamma$ -null sets are precisely the Lebesgue null sets.

There are spaces in which  $\sigma$ -porous sets are  $\Gamma$ -null. Among the classical Banach spaces, the space  $c_0$  is such an example, see [1]. In those cases, the above-quoted statement gives a nice quantitative result saying that a Lipschitz function is Fréchet differentiable except for a  $\Gamma$ -null set. Unfortunately, the spaces  $\ell_p$ ,  $1 < p < \infty$ , do not belong to this class. The reason for that is an example published in [3]. It shows, though in an implicit way and with the help of the mean-value theorem from [1], that every space  $\ell_p$ ,  $1 < p < \infty$ , necessarily contains a  $\sigma$ -porous set which is not  $\Gamma$ -null. Nevertheless, one can still profit from the result about the points of differentiability unless the corresponding  $\sigma$ -porous set is so

big that its complement is  $\Gamma$ -null. A recent example in [2], however, makes clear that such a huge  $\sigma$ -porous set does exist. Namely, *there is a  $\sigma$ -porous set in a separable Hilbert space such that its complement intersects every three dimensional non degenerate  $C^1$  surface in a set of measure zero*. This property is sufficient for a set to have a  $\Gamma$ -null complement. It is interesting to note that if we use two-dimensional surfaces instead of three-dimensional, the statement is false. For another result in [2], it was shown that given a porous set  $M$  in a separable Hilbert space, almost all points of a typical two-dimensional surface do not belong to  $M$ .

One of the most restrictive notions of smallness is the so-called unrectifiability. A set is called 1-purely unrectifiable if the intersection with any  $C^1$  curve is of  $\mathcal{H}^1$ -measure zero. The space  $\ell_1$  contains a  $\sigma$ -porous set whose complement is 1-purely unrectifiable, see [2]. From what was said above, it is clear that such a big  $\sigma$ -porous set cannot occur in every (separable) Banach space, for example not in  $c_0$ . However, if we relax a bit the requirement for porosity, then it is possible to construct in every Banach space the “generalized”  $\sigma$ -porous set with 1-purely unrectifiable complement, see Theorem 2.2 below. An analogous result for standard  $\sigma$ -porous set holds true provided that we use only lines instead of all  $C^1$  curves, see [3]: *in every separable Banach space, there is a  $\sigma$ -porous set the complement of which is of measure zero on every line*.

Although the generalized porous set does not seem to have a direct connection to differentiation of Lipschitz functions, we believe that the result presented in the next section may be of some interest. At least it answers the question which naturally appeared in this context: how big can a  $g$ -porous set be?

## 2. Main result

We start with the notion of a  $g$ -porous set. This concept is a direct generalization of the standard porosity. It can be found also in the above-mentioned survey paper [4] together with the more detailed treatment of various other notions of porous sets. In what follows, we denote by  $B(x, r)$  the ball with centre  $x$  and radius  $r$ .

**Definition 2.1.** Let  $g : [0, \infty) \rightarrow [0, \infty)$  be a continuous increasing function, and let  $g(0) = 0$ . A set  $M \subset X$  in a metric space  $X$  is called  $g$ -porous if for any  $x \in M$ ,

$$\limsup_{s \rightarrow 0_+} \frac{g(\gamma(x, s, M))}{s} > 0, \quad (2.1)$$

where  $\gamma(x, s, M) = \sup \{r \geq 0 \mid B(y, r) \cap M = \emptyset, \text{dist}(x, y) \leq s\}$ .

A set which is a countable union of  $g$ -porous sets is called a  $\sigma$ - $g$ -porous set.

Every  $g$ -porous set is a nowhere dense set and, consequently,  $\sigma$ - $g$ -porous set is a set of the 1st category. If  $g(t) = t$ , then  $g$ -porosity becomes the standard porosity. Notice also that if  $g'_+(0)$  is finite and nonzero (or, more generally,  $0 < \underline{D}_+g(0) \leq \overline{D}_+g(0) < \infty$ ), then a  $g$ -porous set is again a porous set only. If we allow  $g'_+(0) = +\infty$ , then we get a type of sets which are in general bigger than porous sets. The next theorem deals exactly with such types of  $g$ -porous sets.

The symbols  $\mathcal{H}^1$  and  $\mathcal{L}^1$  denote one-dimensional Hausdorff and Lebesgue measures, respectively.

**THEOREM 2.2.** *Let  $X$  be a Banach space and let  $g : [0, \infty) \rightarrow [0, \infty)$  be a continuous, increasing function with  $g(0) = 0$  and*

$$\limsup_{s \rightarrow 0_+} \frac{g(s)}{s} = \infty. \quad (2.2)$$

*Then there is a  $\sigma$ - $g$ -porous set  $M \subset X$  such that the complement  $X \setminus M$  meets every  $C^1$  curve in a set of  $\mathcal{H}^1$ -measure zero. In short,  $X \setminus M$  is purely 1-unrectifiable.*

**Remark 2.3.** It is useful to realize the following equivalent reformulation of the conclusion of Theorem 2.2. If  $\varphi : \mathbb{R} \rightarrow X$  is any  $C^1$  curve, then

$$\mathcal{H}^1(\varphi(\mathbb{R}) \cap (X \setminus M)) = 0 \quad (2.3)$$

if and only if

$$\mathcal{L}^1\{t \in \mathbb{R} \mid \varphi(t) \notin M, \varphi'(t) \neq 0\} = 0. \quad (2.4)$$

This is an immediate consequence of the formula for the length of a curve as follows:

$$\begin{aligned} \mathcal{H}^1(\varphi(\mathbb{R}) \cap (X \setminus M)) &= \int_{\{t \mid \varphi(t) \notin M\}} \|\varphi'(t)\| d\mathcal{L}^1(t) \\ &= \int_{\{t \mid \varphi(t) \notin M, \varphi'(t) \neq 0\}} \|\varphi'(t)\| d\mathcal{L}^1(t). \end{aligned} \quad (2.5)$$

We prove first a simple lemma.

**LEMMA 2.4.** *Let  $\varphi : \mathbb{R} \rightarrow X$  and let  $v \in X$ ,  $v \neq 0$ , be such that  $\varphi(t) - vt$  is a  $K$ -Lipschitz mapping with  $0 \leq K < \|v\|$ .*

(i) *For any  $s, t \in \mathbb{R}$ ,*

$$(\|v\| - K)|s - t| \leq \|\varphi(s) - \varphi(t)\| \leq (\|v\| + K)|s - t|. \quad (2.6)$$

(ii) *If  $P \subset X$  is a nonempty set, then*

$$\mathcal{H}^1(\varphi(\mathbb{R}) \cap P) \leq \frac{\|v\| + K}{\|v\| - K} \text{diam } P. \quad (2.7)$$

**Remark 2.5.** The assumption that  $\varphi(t) - vt$  is a  $K$ -Lipschitz mapping is a weakening of a more geometrically apparent condition which one can use in the case of  $C^1$  mapping, namely,  $\|\varphi'(t) - v\| \leq K$ . Indeed, let  $s, t \in \mathbb{R}$ ,  $s \leq t$ , be arbitrary. Then the mean-value theorem gives

$$\begin{aligned} \|\varphi(s) - vs - (\varphi(t) - vt)\| &\leq \sup\{\|\varphi'(\theta) - v\| \mid \theta \in (s, t)\} |s - t| \\ &\leq K|s - t|. \end{aligned} \quad (2.8)$$

*Proof.* (i) Since  $\varphi(t) - vt$  is  $K$ -Lipschitz, we have

$$\begin{aligned} \|\varphi(s) - \varphi(t)\| &\leq \|\varphi(s) - vs - \varphi(t) - vt\| + \|vs - vt\| \\ &\leq K|s - t| + \|v\||s - t| = (\|v\| + K)|s - t|. \end{aligned} \quad (2.9)$$

Similarly, we estimate the expression  $\|\varphi(s) - \varphi(t)\|$  from below, which gives (i).

(ii) If  $\text{diam } P = \infty$ , the statement is clear. Assume  $\text{diam } P < \infty$ . We denote

$$\begin{aligned} t_0 &= \inf \{t \in \mathbb{R} \mid \varphi(t) \in P\}, \\ t_1 &= \sup \{t \in \mathbb{R} \mid \varphi(t) \in P\}. \end{aligned} \quad (2.10)$$

From already proved part (i), we infer that both  $t_0$  and  $t_1$  are finite and that

$$(\|v\| - K)|t_1 - t_0| \leq \|\varphi(t_1) - \varphi(t_0)\| \leq \text{diam } P. \quad (2.11)$$

Hence, using also the fact that  $\mathcal{H}^1(\varphi([t_0, t_1])) \leq (\|v\| + K)|t_1 - t_0|$ , we obtain

$$\begin{aligned} \mathcal{H}^1(\varphi(\mathbb{R}) \cap P) &= \mathcal{H}^1(\varphi([t_0, t_1]) \cap P) \\ &\leq (\|v\| + K)|t_1 - t_0| \\ &\leq \frac{\|v\| + K}{\|v\| - K} \text{diam } P. \end{aligned} \quad (2.12)$$

□

*Proof of Theorem 2.2.* Consider the family  $\mathcal{D}$  of all 1-discrete sets in  $X$ . (We recall that a set  $D$  is called  $\lambda$ -discrete if  $\|d_1 - d_2\| \geq \lambda$  whenever  $d_1, d_2$  are two distinct elements in  $D$ .) By Zorn's lemma,  $\mathcal{D}$  contains a maximal element  $D$ . Further, the assumption (2.2) implies that there is a sequence  $(s_n) \subset \mathbb{R}$ ,  $s_n \searrow 0$ , such that

$$r_n \equiv \frac{s_n}{g(s_n)} \searrow 0, \quad \sum_{n=1}^{\infty} r_n < \infty. \quad (2.13)$$

Denote

$$\mathcal{B}_n = \bigcup \{B(d, r_n) \mid d \in D\}, \quad M_n = X \setminus \bigcup_{k \geq n} g(s_k) \mathcal{B}_k. \quad (2.14)$$

We show that  $M_n$  is  $g$ -porous. Let  $x \in M_n$ , and let  $k \geq n$ . By the maximality of  $D$ , there is  $d \in D$  such that

$$\left\| \frac{x}{g(s_k)} - d \right\| < 1, \quad \text{i.e.,} \quad \|x - g(s_k)d\| < g(s_k). \quad (2.15)$$

The ball  $B(g(s_k)d, s_k) = g(s_k)B(d, r_k) \subset g(s_k)\mathcal{B}_k$  is obviously contained in the complement of  $M_n$ . Hence

$$\gamma(x, g(s_k), M_n) \geq s_k. \quad (2.16)$$

The ratio of the  $g$ -porosity can be now estimated from below:

$$\begin{aligned} \limsup_{s \rightarrow 0_+} \frac{g(\gamma(x, s, M_n))}{s} &\geq \limsup_{k \rightarrow \infty} \frac{g(\gamma(x, g(s_k), M_n))}{g(s_k)} \\ &\geq \limsup_{k \rightarrow \infty} \frac{g(s_k)}{g(s_k)} = 1, \end{aligned} \quad (2.17)$$

and the  $g$ -porosity of  $M_n$  is proved.

We put  $M = \bigcup_n M_n$ . Clearly,  $M$  is a  $\sigma$ - $g$ -porous set.

Let  $\varphi: \mathbb{R} \rightarrow X$  be any  $C^1$  curve. In view of Remark 2.3, we finish the proof by showing that the set

$$\{t \in \mathbb{R} \mid \varphi(t) \notin M, \varphi'(t) \neq 0\} \quad (2.18)$$

has  $\mathcal{L}^1$ -measure zero. To this end, let  $t_0$  be an arbitrary point from this set and let  $v = \varphi'(t_0)$ . Then  $v$  is a nonzero vector. By continuity of  $\varphi'$ , there is a compact interval  $I$  containing  $t_0$  in the interior and a number  $0 \leq K < \|v\|$  verifying that

$$\|\varphi'(t) - v\| < K, \quad t \in I. \quad (2.19)$$

Since

$$\begin{aligned} \varphi(I) \cap (X \setminus M) &= \varphi(I) \cap \bigcap_{n=1}^{\infty} (X \setminus M_n) \\ &= \varphi(I) \cap \bigcap_{n=1}^{\infty} \left( \bigcup_{k \geq n} g(s_k) \mathcal{B}_k \right) \\ &\subset \varphi(I) \cap \left( \bigcup_{k \geq n} g(s_k) \mathcal{B}_k \right) \\ &= \bigcup_{k \geq n} (\varphi(I) \cap g(s_k) \mathcal{B}_k) \end{aligned} \quad (2.20)$$

for any  $n \in \mathbb{N}$ , it follows that

$$\mathcal{H}^1(\varphi(I) \cap (X \setminus M)) \leq \sum_{k \geq n} \mathcal{H}^1(\varphi(I) \cap g(s_k) \mathcal{B}_k). \quad (2.21)$$

By Remark 2.5, the mapping  $\varphi(t) - vt$  is  $K$ -Lipschitz on  $I$ . Thus we can apply Lemma 2.4(ii) to each ball  $B(g(s_k)d, s_k)$  which meets the arc  $\varphi(I)$  and we get

$$\mathcal{H}^1(\varphi(I) \cap B(g(s_k)d, s_k)) \leq \frac{\|v\| + K}{\|v\| - K} 2s_k. \quad (2.22)$$

Now let  $n \in \mathbb{N}$  be large enough to guarantee that  $1 \geq g(s_k) \geq 3s_k$  for all  $k \geq n$ . (This is possible due to (2.13).) Let  $k \geq n$  and let  $N_k$  denote the number of balls  $B(g(s_k)d, s_k)$ ,  $d \in D$ , intersecting the arc  $\varphi(I)$ . Since  $D$  is 1-discrete, the distance between any two distinct

points from  $g(s_k)D$  is at least  $g(s_k)$ . Hence the part of the arc  $\varphi(I)$  between two consecutive balls has measure at least  $g(s_k) - 2s_k$ . So we infer that

$$(N_k - 1)(g(s_k) - 2s_k) \leq \mathcal{H}^1(\varphi(I) \setminus g(s_k)\mathcal{B}_k). \quad (2.23)$$

The mapping  $\varphi$  itself is  $(\|v\| + K)$ -Lipschitz on  $I$  by Lemma 2.4(i). So

$$\mathcal{H}^1(\varphi(I) \setminus g(s_k)\mathcal{B}_k) \leq \mathcal{H}^1(\varphi(I)) \leq (\|v\| + K)\mathcal{L}^1(I). \quad (2.24)$$

Combining (2.23) and (2.24), we obtain an upper estimate for the number  $N_k$  of balls meeting  $\varphi(I)$ :

$$N_k \leq 1 + \frac{(\|v\| + K)\mathcal{L}^1(I)}{g(s_k) - 2s_k}. \quad (2.25)$$

With the help of (2.22), we get

$$\begin{aligned} \mathcal{H}^1(\varphi(I) \cap g(s_k)\mathcal{B}_k) &= \sum_{d \in D} \mathcal{H}^1(\varphi(I) \cap B(g(s_k)d, s_k)) \\ &\leq N_k \frac{\|v\| + K}{\|v\| - K} 2s_k \\ &\leq \left[ 1 + \frac{(\|v\| + K)\mathcal{L}^1(I)}{g(s_k) - 2s_k} \right] \frac{\|v\| + K}{\|v\| - K} 2s_k \\ &\leq \left[ 1 + \frac{(\|v\| + K)\mathcal{L}^1(I)}{2/3} \right] \frac{\|v\| + K}{\|v\| - K} \frac{2s_k}{g(s_k)} \leq Cr_k, \end{aligned} \quad (2.26)$$

where the constant  $C$  depends only on  $\mathcal{L}^1(I)$  and  $K$ . Consequently, in view of (2.21),

$$\mathcal{H}^1(\varphi(I) \cap (X \setminus M)) \leq C \sum_{k \geq n} r_k. \quad (2.27)$$

This is true for all  $n$  sufficiently big. So we conclude that the interval  $I$  possesses the property

$$\mathcal{H}^1(\varphi(I) \cap (X \setminus M)) = 0, \quad \text{i.e.,} \quad \mathcal{L}^1\{t \in I \mid \varphi(t) \notin M, \varphi'(t) \neq 0\} = 0. \quad (2.28)$$

By separability of  $\mathbb{R}$ , the set  $\{t \in \mathbb{R} \mid \varphi(t) \notin M, \varphi'(t) \neq 0\}$  can be covered by countably many of such intervals  $I$ . This fact, together with (2.28), finishes the proof.  $\square$

### Acknowledgment

The author was supported by Grants GAČR 201/04/0090 and MSM 210000010.

## References

- [1] J. Lindenstrauss and D. Preiss, *On Fréchet differentiability of Lipschitz maps between Banach spaces*, Ann. of Math. (2) **157** (2003), no. 1, 257–288.
- [2] J. Lindenstrauss, D. Preiss, and J. Tišer, *Avoiding  $\sigma$ -porous sets*, in preparation.
- [3] D. Preiss and J. Tišer, *Two unexpected examples concerning differentiability of Lipschitz functions on Banach spaces*, Geometric Aspects of Functional Analysis (Israel, 1992–1994), Oper. Theory Adv. Appl., vol. 77, Birkhäuser, Basel, 1995, pp. 219–238.
- [4] L. Zajíček, *Porosity and  $\sigma$ -porosity*, Real Anal. Exchange **13** (1987/1988), no. 2, 314–350.

Jaroslav Tišer: Department of Mathematics, Faculty of Electrical Engineering, Czech Technical University in Prague, Žitkova 4, 166 27 Praha 6, Czech Republic

*E-mail address:* tiser@math.feld.cvut.cz

# Differential Equations & Nonlinear Mechanics

*An Open Access Journal*

## Editor-in-Chief

K. Vajravelu  
USA

## Associate Editors

N. Bellomo  
*Italy*

J. L. Bona  
USA

J. R. Cannon  
USA

S.-N. Chow  
USA

B. S. Dandapat  
*India*

E. DiBenedetto  
USA

R. Finn  
USA

R. L. Fosdick  
USA

J. Frehse  
*Germany*

A. Friedman  
USA

R. Grimshaw  
*UK*

J. Malek  
*Czech Republic*

J. T. Oden  
USA

R. Quintanilla  
*Spain*

K. R. Rajagopal  
USA

G. Saccomandi  
*Italy*

Y. Shibata  
*Japan*

Ivar Stakgold  
USA

Swaroop Darbha  
USA

A. Tani  
*Japan*

S. Turek  
*Germany*

A. Wineman  
USA

**Website:** <http://www.hindawi.com/journals/denm/>

## Aims and Scope

Differential equations play a central role in describing natural phenomena as well as the complex processes that arise from science and technology. Differential Equations & Nonlinear Mechanics (DENM) will provide a forum for the modeling and analysis of nonlinear phenomena. One of the principal aims of the journal is to promote cross-fertilization between the various subdisciplines of the sciences: physics, chemistry, and biology, as well as various branches of engineering and the medical sciences.

Special efforts will be made to process the papers in a speedy and fair fashion to simultaneously ensure quality and timely publication.

DENM will publish original research papers that are devoted to modeling, analysis, and computational techniques. In addition to original full-length papers, DENM will also publish authoritative and informative review articles devoted to various aspects of ordinary and partial differential equations and their applications to sciences, engineering, and medicine.

## Open Access Support

The Open Access movement is a relatively recent development in academic publishing. It proposes a new business model for academic publishing that enables immediate, worldwide, barrier-free, open access to the full text of research articles for the best interests of the scientific community. All interested readers can read, download, and/or print any Open Access articles without requiring a subscription to the journal in which these articles are published.

In this Open Access model, the publication cost should be covered by the author's institution or research funds. These Open Access charges replace subscription charges and allow the publishers to give the published material away for free to all interested online visitors.

## Instructions for Authors

Original articles are invited and should be submitted through the DENM manuscript tracking system at <http://www.mstracking.com/denm/>. Only pdf files are accepted. If, for some reason, submission through the manuscript tracking system is not possible, you can contact [denm.support@hindawi.com](mailto:denm.support@hindawi.com).

---

## Hindawi Publishing Corporation

410 Park Avenue, 15th Floor, #287 pmb, New York, NY 10022, USA

**HINDAWI**

<http://www.hindawi.com/journals/denm/>